

11/28/00



jc690 U.S. PTO

11-29-00

Docket No. 0575/57006/JPW/JML

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Honorable Assistant Commissioner for Patents
Washington, D.C. 20231
BOX: PATENT APPLICATION
S I R:

November 28, 2000

jc813 U.S. PTO

09/724028



Transmitted herewith for filing are the specification and claims of the patent application of:

Paul B. Fisher and Dong-chul Kang for
Inventor(s)

METHOD FOR FULL-LENGTH cDNA CLONING USING DEGENERATE STEM LOOP ANNEALING PRIMERS
Title of Invention

Also enclosed are:

☒ 7 sheet(s) of ☐ informal ☒ formal drawings.

☒ Oath or declaration of Applicant(s). (Unsigned)

☒ A power of attorney (Unsigned)

☐ An assignment of the invention to _____

☐ A Preliminary Amendment

☐ A verified statement to establish small entity status under 37 C.F.R. §1.9 and §1.27.

The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

	NUMBER FILED		NUMBER EXTRA*		RATE		FEE	
					SMALL ENTITY	OTHER ENTITY	SMALL ENTITY	OTHER ENTITY
Total Claims	18 -20	=	0	X	\$ 9.00	\$18.00	= \$ 0	\$
Independent Claims	4 -3	=	1	X	\$40.00	\$80.00	= \$ 40.00	\$
Multiple Dependent Claims Presented: <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No					\$135.00	\$270.00	= \$ 135.00	\$
*If the different in Col. 1 is less than zero, enter "0" in Col. 2					BASIC FEE		\$ 355.00	\$ 710
					TOTAL FEE		\$ 530.00	\$

X A check in the amount of \$ 530.00 to cover the filing fee.

 Please charge Deposit Account No. in the amount of \$.

X The Commissioner is hereby authorized to charge any additional fees which may be required in connection with the following or credit any over-payment to Account No. 03-3125:

X Filing fees under 37 C.F.R. §1.16.

X Patent application processing fees under 37 C.F.R. §1.17.

 The issue fee set in 37 C.F.R. §1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. §1.311(b).

 Three copies of this sheet are enclosed.

 A certified copy of previously filed foreign application No. filed in on .

Applicant(s) hereby claim priority based upon this aforementioned foreign application under 35 U.S.C. §119.

X Other (identify) One extra set of formal drawings and an Express Mail Certificate of Mailing bearing label number EJ 900 852 073 US.

Respectfully submitted,

John P. White

John P. White
Registration No. 28,678
Jane M. Love
Registration No. 42,812
Attorney for Applicants
Cooper & Dunham, LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Paul B. Fisher and Dong-chul Kang
Application No.: Not Yet Known
Filing date : Herewith
For : METHOD FOR FULL-LENGTH cDNA CLONING USING
DEGENERATE STEM-LOOP ANNEALING PRIMERS

1185 Avenue of the Americas
New York, New York 10036
November 28, 2000

Assistant Commissioner for Patents
Washington, D.C. 20231
BOX: Patent Application

SIR:

EXPRESS MAIL CERTIFICATE OF
MAILING FOR ABOVE-IDENTIFIED APPLICATION

"Express Mail" mailing label number: EJ 900 852 073 US
Date of Deposit: November 28, 2000
I hereby certify that this paper or fee is being deposited with
the United States Postal Service "Express Mail Post Office to
Addressee" service under 37 C.F.R. §1.10 on the date indicated
above and is addressed to the Assistant Commissioner for Patents
Box Patent Application, Washington, D.C. 20231.

Printed Name: John P. White

Respectfully submitted,

Jane M. Love

John P. White
Registration No. 28,678
Jane M. Love
Registration No. 42,812
Attorney for Applicant(s)
Cooper & Dunham, LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

008871 3226450

*Application
for
United States Letters Patent*

09724028-112800

To all whom it may concern:

Be it known that **Paul B. Fisher and Dong-Chul Kang**

have invented certain new and useful improvements in

Method for Full-Length cDNA Cloning Using Degenerate Stem-Loop Annealing Primers

of which the following is a full, clear and exact description.

5

10

15

25

The human genome is estimated to contain 100,000 genes, the expressions of which define the functionality of a cell (1). Current technological advances, including large-scale DNA

sequencing, efficient library construction and manipulation and PCR-based gene expression monitoring, have resulted in the identification of more than 87,000 unique expression sequence tags (ESTs) in diverse cell types and under various physiological conditions (1). Approximately 12 % of the ESTs have significant homology with previously identified genes and the remainder require further investigation to define their identity and biological relevance (1). However, ESTs, short stretches of expressed genes, can only provide limited information as to the identity and biological role of specific genes. A more thorough analysis of the ESTs requires a determination of the full protein coding sequences for these expressed genes.

Several approaches are routinely used to obtain cDNAs containing protein-encoding sequences from ESTs. These include, library screening (2) and the PCR-based rapid amplification of cDNA ends (RACE) strategy (3). A less frequently employed scheme, exon trapping is also amenable to cDNA cloning from genomic fragments (4,5).

A number of cDNA libraries from diverse sources are commercially available. This can in specific instances reduce the burden of producing cDNA libraries that are required for screening for cDNAs. However, even with well-constructed cDNA libraries, several rounds of screening and verification are often required to obtain even a single complete cDNA (2). This process is laborious and can require months of intensive effort. What exacerbates the

situation is that cDNA library screening occasionally results in incomplete cDNAs lacking full protein coding information. This occurs primarily because of premature termination of reverse transcription and the self-priming procedure during second strand cDNA synthesis (2, 6, 7). Additionally, obtaining cDNA of low abundance mRNA is rarely achievable unless the cDNA library is high titer and minimally amplified (2). In these contexts, the current approach of cDNA library screening to obtain full protein coding sequence is often costly, laborious and inefficient.

09724029-1330

Summary of the Invention

The present invention provides a method for isolating a double-stranded cDNA having a nucleotide sequence of a complete open reading frame which comprises: (A) admixing (i) an isolated single-stranded cDNA, (ii) a first primer capable of forming a stem-loop structure, comprising (a) at the 3' end of the primer, a first random sequence, linked to (b) a second sequence, linked to (c) a third sequence which forms a loop structure, linked to (d) a fourth sequence, at the 5' end of the first primer, which is complementary to the second sequence, under hybridization conditions sufficient for annealing the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and (iii) a polymerase; (B) incubating the mixture from step (A) under suitable conditions for DNA synthesis; and (C) performing a polymerase chain reaction by admixing (i) an aliquot of the mixture from (B), (ii) a second primer which specifically binds to the single-stranded cDNA, (iii) a third primer which comprises (a) a fifth sequence identical to the third sequence of the first primer, linked to (b) a sixth sequence identical to a portion of the second sequence of the first primer, and (iv) a polymerase, under conditions suitable for a polymerase chain reaction so as to produce a double-stranded cDNA reaction product, thereby isolating the cDNA having the sequence of the complete open reading frame.

Brief Description of the Figures

Figures 1A-C: Schematic of the C-ORF procedure and the primers used in this approach. **Fig 1A.** In C-ORF, specifically designed primers anneal to the 3' end of the first strand cDNA. Extended cDNA from this primer can be amplified by PCR with a gene specific primer (GSP) and a universal anchor primer containing the annealing primer sequence. This method dispenses with rate limiting steps of conventional RACE, such as homopolymer tailing or single strand ligation. **Fig 1B.** Primers (D-SLAP, D-CLAP1 and D-CLAP2) used in the 5' C-ORF reaction. The stem-and-loop structure of these primers is designed to prevent, by steric hindrance, degenerate regions from annealing to internal sites, thereby insuring efficient extension of the gene product. D-CLAP1 contains additional GGG sequences to enhance annealing to the C-homopolymer tail generated by the TdT-like activity of reverse transcriptase (RT). **Fig 1C.** In the T-SLAP reagent, dT replaces the random sequences used for the reverse transcription reaction as a primer.

Figures 2A-2E. C-ORF cloning of the novel gene *mda-5*. **Fig 2A.** *Mda-5* C-ORF products (12 μ l nested PCR) with reverse transcription reactions at specified temperature are resolved in a 1% agarose gel containing EtBr. **Fig 2B.** C-ORF products of *mda-5* performed with different amounts of the D-SLAP reagent and the anchor primer in primary PCR for second strand cDNA synthesis. **Fig 2C.** An autoradiogram of Southern blot hybridization of C-ORF products of *mda-5*

resolved in (B) with ^{32}P -labeled nested primers. **Fig 2D.**
An autoradiogram of Northern blot hybridization of RNA
samples prepared from HO-1 human melanoma cells either left
alone (Con) or treated with 2,000 U/ml IFN- β plus 10 ng/ml
5 mezerein. Left panel was probed with ^{32}P -labeled *mda-5* EST
(0.4 kb) previously cloned by library screening. The right
panel was probed with a ^{32}P -labeled 1.8-kb *mda-5* C-ORF
product. **Fig 2E.** RT-PCR analysis using sequence
information derived from the *mda-5* C-ORF product. RT-PCR
10 was performed using 2 μl of the reverse transcription
reaction with the specified primer, either R2S8 or R4S8, and
12 μl of the reaction was run on the gel.

Figure 3. Applications of the C-ORF protocol for
15 identification of the complete ORF of ISG-56, *mda-9* and *mda-5*.
C-ORF products of ISG-56, *mda-9* and *mda-5* were resolved
in 1 % agarose gel (lanes 3, 6 and 9). The C-ORF products
are shown in comparison with the RT-PCR products of each
gene using a common 3' nested primer and a 5' primer from
20 reported gene sequence (lanes 2, 5 and 8). Nested PCR of C-
ORF with only an anchor primer (lanes 4, 7 and 10)
distinguishes gene specific C-ORF products from RT-PCR
artifacts.

25 **Figures 4A-4B.** C-ORF protocol with the D-CLAP1 and D-CLAP2
reagents for 5' cDNA end cloning. **Fig 4A.** C-ORF products
obtained using the D-CLAP1 reagent for the specified genes
were separated in a 1% agarose gel. Authentic bands of the
appropriate target size are marked with dots. Duration of

5

10

15

Detailed Description of the Invention

The present invention provides for a method for isolating a double-stranded cDNA having a nucleotide sequence of a complete open reading frame which comprises: A) admixing (i) an isolated single-stranded cDNA, (ii) a first primer capable of forming a stem-loop structure, comprising (a) at the 3' end of the primer, a first random sequence, linked to (b) a second sequence, linked to (c) a third sequence which forms a loop structure, linked to (d) a fourth sequence, at the 5' end of the first primer, which is complementary to the second sequence, under hybridization conditions sufficient for annealing the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and (iii) a polymerase; B) incubating the mixture from step (A) under suitable conditions for DNA synthesis; and C) performing a polymerase chain reaction by admixing (i) an aliquot of the mixture from (B), (ii) a second primer which specifically binds to the single-stranded cDNA, (iii) a third primer which comprises (a) a fifth sequence identical to the third sequence of the first primer, linked to (b) a sixth sequence identical to a portion of the second sequence of the first primer, and (iv) a polymerase under conditions suitable for a polymerase chain reaction so as to produce a double-stranded cDNA reaction product, thereby isolating the cDNA having the sequence of the complete open reading frame.

In one embodiment of the invention, the single-stranded DNA is a 5' portion of a cDNA reverse transcribed from an mRNA.

In another embodiment of the invention, the first primer has the sequence 3'-

NNNNNNNNNNNNNCAGAGCTCAAATTTGTGATCAGCTGGTCTTTCACAAATTTGAGCTC
TG-5' (D-SLAP).

- 5 In another embodiment of the invention, the first primer has the sequence

3'NNNNNNNNNNNGGGAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGTAGAAT
ACATCTTGAGCTAT-5' (D-CLAP1).

- 10 In a further embodiment of the invention, the first primer has the sequence

3'NNNNNNNNNNNNNNNAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGTAGAAT
ACATCTTGAGCTAT(D-CLAP2).

- 15 In another embodiment of the invention, the first primer comprises an inosine nucleotide.

In a further embodiment of the invention, the loop structure is a simple loop structure, or a cloverleaf loop structure.

20

- The present invention provides for a method for generating a cDNA library which comprises: A) admixing (i) a population of single-stranded cDNA molecules which were reverse transcribed with an oligo-dT sequence linked to a defined
25 nucleotide sequence, (ii) a first primer capable of forming a stem-loop structure, comprising (a) at the 3' end of the primer, a first random sequence linked to (b) a second sequence, linked to (c) a third sequence which forms a loop structure, linked to (d) a fourth sequence, at the 5' end of

09724038-12800

the first primer, which is complementary to the second sequence, under hybridization conditions sufficient for annealing the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and
5 (iii) a polymerase; B) incubating the mixture from step (A) under suitable conditions for DNA synthesis by the polymerase; and C) performing a polymerase chain reaction by admixing (i) an aliquot of the mixture from (B), (ii) a second primer which has the identical sequence as the
10 defined nucleotide sequence of the primer in (A)(i), (iii) a third primer which comprises (a) a fifth sequence identical to the third sequence of the first primer, linked to (b) a sixth sequence identical to a portion of the second sequence of the first primer, and (iv) a polymerase under
15 conditions suitable for a polymerase chain reaction so as to produce double-stranded cDNA reaction products thereby generating a cDNA library.

In one embodiment, the single-stranded DNA is a cDNA reverse
20 transcribed from an mRNA.

In another embodiment of the invention, the first primer has the sequence 3'-
NNNNNNNNNNNNNCAGAGCTCAAATTTGTGATCAGCTGGTCTTTCACAAATTTGAGCTC
25 TG-5' (D-SLAP).

In another embodiment of the invention, the first primer has the sequence
3' NNNNNNNNNNNGGAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGTAGAAT

ACATCTTGAGCTAT-5' (D-CLAP1).

In another embodiment of the invention, the first primer has the sequence

5 3'NNNNNNNNNNNNNAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGTAGAAT
ACATCTTGAGCTAT(D-CLAP2).

In another embodiment of the invention, the first primer comprises an inosine nucleotide.

10

In another embodiment of the invention, the loop structure is a simple loop structure, or a cloverleaf loop structure.

The present invention provides for a kit for the generation
15 of a complete open reading frame double-stranded cDNA of
interest which comprises: (i) a first primer capable of
forming a stem-loop structure, comprising (a) at the 3' end
of the primer, a first random sequence linked to (b) a
second sequence, linked to (c) a third sequence which forms
20 a loop structure, linked to (d) a fourth sequence, at the 5'
end of the first primer, which is complementary to the
second sequence, and (ii) a second primer which comprises
(a) a fifth sequence identical to the third sequence of the
first primer, linked to (b) a sixth sequence identical to a
25 portion of the second sequence of the first primer.

The present invention also provides for a method for
isolating a double-stranded cDNA having a nucleotide
sequence of a complete open reading frame which comprises:

09724028.1 32042/60

00937-824260

(a) admixing (i) a biological sample containing mRNA, (ii) a primer which forms a stem-loop structure, comprising: (a) a poly-T sequence at the 3' end of the primer linked to (b) a first random sequence linked to (c) a second sequence which forms a loop structure linked to (d) a third sequence at the 5' end of the primer which is complementary to the first sequence, and (iii) a reverse transcriptase, under hybridization conditions sufficient for annealing the primer to the mRNA poly-A sequence; (b) incubating the mixture from step (a) under suitable conditions for reverse transcription; (c) performing a polymerase chain reaction with an aliquot of the mixture from step (b) using one gene-specific primer which is pre-defined and one primer which has a sequence identical to at least a portion of the primer sequence of element (ii), thereby isolating the cDNA having the sequence of the complete open reading frame.

In one embodiment of the invention, the primer has the sequence 3' -
TTTTTTTTTTTCAGAGCTCAAATTTGTGATCAGCTGGTCTTTCACAAATTTGAGCTC
TG-5' (T-SLAP).

In addition, the present invention is directed to a method for isolating a double-stranded cDNA having a nucleotide sequence of a complete open reading frame which comprises:
A) admixing (i) an isolated single-stranded cDNA, (ii) a first primer capable of forming a stem-loop structure, comprising (a) at the 3' end of the primer, a first random sequence, linked to (b) a second sequence, linked to (c) a

of the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and (iii) a ligase; B) incubating the mixture from step (A) under suitable conditions for DNA synthesis by the polymerase; and C) performing a polymerase chain reaction by admixing (i) an aliquot of the mixture from (B), (ii) a second primer which has the identical sequence as the defined nucleotide sequence of the primer in (A)(i), (iii) a third primer which comprises (a) a fifth sequence identical to the third sequence of the first primer, linked to (b) a sixth sequence identical to a portion of the second sequence of the first primer, and (iv) a polymerase under conditions suitable for a polymerase chain reaction so as to produce double-stranded cDNA reaction products thereby generating a cDNA library.

15

The following are several applications of the present invention:

1. Cloning the 5' end of a cDNA from an EST.
2. Cloning the 3' end of a cDNA from an EST by performing C-ORF in a reverse transcription reaction.
3. Construction of a cDNA library containing a high proportion of full-length cDNAs.
4. Genomic cloning, both upstream and downstream regions of known sequences.
5. Capture nucleic acid of specific sequence for purification and diagnostic purposes.
6. Gene inactivation by inhibiting mRNA entry onto the ribosome for translation.
7. Cloning family of genes, such as kinases.

9. Chromosomal mapping.

5

15

20

25

libraries and/or the PCR-based 5' RACE approach. Screening
cDNA libraries is very labor intensive and often yields
incomplete cDNA, which necessitates the use of additional 5'
cloning methods such as 5' RACE. To reach the 5' end of a
5 cDNA two PCR-based methods have been used, 5' RACE employing
terminal deoxytransferase to prime the 3' end of the first
strand cDNA and linker ligation using RNA ligase. However,
these methods are limited in effectiveness because of the
low efficiency of the priming procedure. We describe a
10 novel way of priming the 3' end of first strand cDNA that
allows the second strand of cDNA to be synthesized
preferentially from the 3' end of the first strand of cDNA.

Degenerate stem and loop cDNA end annealing primer (D-SLAP)
15 methods were designed to preferentially anneal to the 3' end
of the first strand of cDNA. The primer consists of two
functional components: 1) a stem and a loop (stem-and-loop)
component and 2) the annealing of degenerate nucleotides.
The stem-and-loop structure sterically hinders degenerate
20 oligonucleotides from annealing to the middle of a cDNA and
promotes preferential annealing to the end of the first
strand of cDNA. Conceptually, the stem-and-loop of the D-
SLAP method should form molecular complexes at higher
temperatures preventing the binding of degenerate
25 oligonucleotides. This could occur because the longer
region of potential base pairing in the stem-and-loop region
should facilitate intramolecular annealing over degenerate
oligonucleotide binding. Alternatively, the longer region
of base pairing in the stem-and-loop and the intramolecular

annealing reaction mechanics may facilitate formation of stem-and-loop structure prior to degenerate oligonucleotide annealing.

5 Full-length cloning of a 5' sequence of cDNA using the D-SLAP method was performed as follows. Total RNA (2 μ g) treated with RNase-free DNase was extracted and reverse transcribed with a gene specific primer using MMLV RT (Superscript RT II from Gibco-BRL) as in the 5' RACE
10 protocol (Gibco-BRL) at 48°C and inactivated by incubation at 85°C for 5 minutes. After treatment with a mixture of RNase H and RNase I for 30 minutes at 37°C, first strand cDNA was purified with GlassMax® (Gibco-BRL) and the second strand of cDNA was synthesized using D-SLAP method.
15 Purified first strand cDNA was mixed with 1 pmole of D-SLAP reactants and KlenTaq reaction buffer, and heated at 95°C for 1.5 minutes and slowly cooled down to 45°C. KlenTaq (0.25 μ l, Clontech) and dNTP (0.5 μ l of 10mM, Gibco-BRL) were added to the 45°C equilibrated reaction mixture and
20 subsequently second strand cDNA synthesis was performed by incubation at 68°C for 30 minutes. A PCR reaction was set up with 5 μ l of the cDNA synthesis reaction mixture, 5' anchor primer designed from the loop area of D-SLAP and a gene specific primer. Secondary PCR was performed with 0.5
25 μ l of the primary PCR product, the same 5' anchor primer (for nested PCR primers) and a nested gene specific primer.

D-SLAP second strand synthesis was applied to clone a full-length *mda-5* (melanoma differentiation associated gene-5)

cancer, cardiovascular and infectious disease states.

One purpose of the present invention is to provide novel technologies for the identification, complete open reading frame cDNA cloning and functional analysis of genes relevant to human diseases. In addition, the present invention also utilizes molecular approaches and high throughput screening procedures to identify small molecules and novel gene products that can directly intervene in disease processes.

The present invention provides for efficient production of the complete open reading frames of cDNAs from partial cDNAs and ESTs. This approach is called the Complete Open Reading Frame (C-ORF) cloning method. This method also offers the ability to generate cDNA libraries which contain a high proportion of full-length cDNAs. A modification of this method is wherein a 3' C-ORF which permits cloning of complete open reading frame cDNAs from partial internal cDNAs missing 3' regions.

This invention also provides for functional gene evaluation. This includes the efficient and stable generation of target cells containing inducible genes for direct functional analysis and the identification of down-stream target genes and biochemical pathways mediating biological responses. These methods include the use of promoters to drive inducible target genes that are more resistant to loss of gene expression after integration and vector constructs that induce tight-regulation of target gene expression.

In addition, this invention also provides for promoter generation and analysis methods which will permit the rapid identification of the promoter region of potentially important genes regulating disease states. The promoters are used as part of a small molecule screening effort, the Rapid Promoter Screening (RPS) approach, to identify potential inhibitors and activators of disease gene transcription. These small molecules will be tested for efficacy in ameliorating specific disease states.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology and recombinant DNA technology which are within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Second Edition (1989); DNA Cloning, Vols. I and II (D. N. Glover ed. 1985); Oligonucleotide Synthesis (M. J. Gait ed. 1984); Nucleic Acid Hybridization (B. D. Hames & S. J. Higgins eds. 1984); Animal Cell Culture (R. K. Freshney ed. 1986); Immobilized Cells and Enzymes (IRL press, 1986); Perbal, B., A Practical Guide to Molecular Cloning (1984); the series, Methods In Enzymology (S. Colowick and N. Kaplan eds., Academic Press, Inc.); and Handbook of Experimental Immunology, Vols. I-IV (D. M. Weir and C. C. Blackwell eds., 1986, Blackwell Scientific Publications).

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references

procedures, calcium phosphate precipitation (Graham, F. L. and Van der Eb, A. J. (1973) Virology 52:456-467), protoplast fusion, liposome-mediated transfer, polybrene-mediated transfection and direct microinjection of the DNA into nuclei. Bacterial cells will generally be transformed using calcium chloride, either alone or in combination with other divalent cations and DMSO (Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Second Edition (1989)). DNA can also be introduced into bacterial cells by electroporation. Methods of introducing exogenous DNA into yeast hosts typically include either the transformation of spheroplasts or transformation of intact yeast cells treated with alkali cations.

Site-specific DNA cleavage is performed by treating with the suitable restriction enzyme (or enzymes) under conditions which are generally understood in the art, and the particulars of which are specified by the manufacturer of these commercially available restriction enzymes (See, e.g. New England Biolabs Product Catalog). In general, about 1 μ g of plasmid or DNA sequences is cleaved by one unit of enzyme in about 20 μ l of buffer solution. Typically, an excess of restriction enzyme is used to insure complete digestion of the DNA substrate. Incubation times of about one hour to two hours at about 37° C. are workable, although variations can be tolerated. After each incubation, protein is removed by extraction with phenol/chloroform, and may be followed by ether extraction, and the nucleic acid recovered from aqueous fractions by precipitation with ethanol. If

Restriction cleaved fragments may be blunt ended by treating with the large fragment of *E. coli* DNA polymerase I (Klenow) in the presence of the four deoxynucleotide triphosphates (dNTPs) using incubation times of about 15 to 25 min at 20°C to 25°C in 50 mM Tris (pH 7.6) 50 mM NaCl, 6 mM MgCl₂, 6 mM DTT and 5-10 μM dNTPs. The Klenow fragment fills in at 5' sticky ends but chews back protruding 3' single strands, even though the four dNTPs are present. If desired, selective repair can be performed by supplying only one of the dNTPs, or with selected dNTPs, within the limitations dictated by the nature of the sticky ends. After treatment with Klenow, the mixture is extracted with phenol/chloroform and ethanol precipitated. Treatment under appropriate conditions with S1 nuclease or Bal-31 results in hydrolysis of any single-stranded portion.

Ligations are performed in 10-50 μ l volumes under the following standard conditions and temperatures using T4 DNA
25 ligase. Ligation protocols are standard (D. Goeddel (ed.)
Gene Expression Technology: Methods in Enzymology (1991)).
In vector construction employing "vector fragments", the
vector fragment is commonly treated with bacterial alkaline
phosphatase (BAP) or calf intestinal alkaline phosphatase

(CIP) in order to remove the 5' phosphate and prevent religation of the vector. Alternatively, re-ligation can be prevented in vectors which have been double digested by additional restriction enzyme digestion of the unwanted
5 fragments.

The Complete Open Reading Frame (C-ORF) Technology: A Simple and Efficient Approach for Obtaining the Entire Protein Coding Region of Genes

10

The ability to analyze a gene's function often requires the identification of the protein-coding region of that gene. Although a number of approaches, including library screening and rapid amplification of cDNA ends (RACE), have been used
15 extensively to identify the complete open reading frame (ORF) of specific cDNA's, these approaches can be inefficient, time consuming and costly. An approach is described, the C-ORF (complete open reading frame) technology, that results in the rapid and efficient
20 identification of protein coding regions of genes in which limited sequence information is available. This scheme was applied successfully, in the majority of cases involving only a single application, in identifying and cloning the complete ORF of genes ranging in size from 1.2 to 8 kb. The
25 C-ORF approach will prove valuable in efforts designed to define the function of a gene in situations where only expressed sequence tags (ESTs) or incomplete cDNA genetic information is available. This strategy offers promise for accelerating the pace of gene discovery and for rapidly

09724038-13300

advancing the fields of functional genomics, proteomics and pharmacogenomics.

5 The human genome is estimated to contain 100,000 genes, the expressions of which define the functionality of a cell (1). Current technological advances, including large-scale DNA sequencing, efficient library construction and manipulation and PCR-based gene expression monitoring, have resulted in the identification of more than 87,000 unique expression
10 sequence tags (ESTs) in diverse cell types and under various physiological conditions (1). Approximately 12 % of the ESTs have significant homology with previously identified genes and the remainder require further investigation to define their identity and biological relevance (1).
15 However, ESTs, short stretches of expressed genes, can only provide limited information as to the identity and biological role of specific genes. A more thorough analysis of the ESTs requires a determination of the full protein coding sequences for these expressed genes.

20 Several approaches are routinely used to obtain cDNAs containing protein-encoding sequences from ESTs. These include, library screening (2) and the PCR-based rapid amplification of cDNA ends (RACE) strategy (3). A less
25 frequently employed scheme, exon trapping is also amenable to cDNA cloning from genomic fragments (4,5).

A number of cDNA libraries from diverse sources are commercially available. This can in specific instances

Recent improvements in cDNA library construction can significantly increase the proportion of full-length cDNAs. These approaches include trapping m⁷GTP-cap in RNA-DNA hybrid (8), ligation of oligonucleotides to de-capped mRNA with T4 RNA ligase (9) and Cap-switch cDNA Library Synthesis Kit utilizing the terminal deoxyribonucleotide transferase (TdT)-like activity of reverse transcriptase (10-12). However, some of these newer protocols involve additional steps including cumbersome chemical or enzymatic reactions (8, 9). Although cDNA library screening can identify mRNA variants and provide more reliable sequence information than PCR, the construction of high-quality cDNA libraries is not

5

10

15

20

25

The application of RACE for obtaining full-length cDNAs, however, is not as simple and straightforward as the theory behind the technique suggests. Problems, such as premature termination of reverse transcription because of the secondary structure of the mRNA, is common to both RACE and cDNA library construction (6). Thermostable RT (14) or the addition of trehalose (21) permits reverse transcription at higher temperatures (~50 to 60°C) and can reduce secondary structure formation. PCR-related problems encountered using the various RACE procedures include:

1. Generation of non-specific products due to the insufficient specificity and priming of the universal primer to intragenic sites.
2. A high rate of false incorporation of bases frequently occurs using Taq polymerase, a problem which can be reduced (3-to-5-fold), but not completely eliminated, by using enzymes with proofreading activity.
3. Amplification biases often resulting in the preferential amplification of shorter products (less than 1 kb).
4. Production of artifacts by amplification of incorrect hybrid molecules resulting from template self-annealing or mispriming (13, 20).

Thus, amplification of shorter fragments than target length probably arises from premature termination of reverse

transcription, intragenic priming by universal primers, and/or the denaturation and annealing kinetics of PCR reactions (22). Consequently, application of the RACE approach for producing complete protein coding sequence of
5 long messages require successive rounds of this procedure, which obviates the time and laborsaving features of this method.

A rate limiting step in the RACE process is the generation
10 of a second universal primer site by an enzymatic reaction of TdT or T4 RNA ligase. However, not only is the TdT reaction inefficient, but also the length of homopolymer tails added by TdT is difficult to control, resulting in heterogeneous PCR products that are troublesome to sequence
15 (13, 15, 18, 23). T4 RNA ligase is also inefficient, especially with longer substrates, and it requires a high concentration of substrate because of its high K_m (millimolar) (24). Furthermore, donor oligonucleotides need to be phosphorylated at their 5' end and deoxygenated at
20 their 3' end to avoid ligation to the 5' end of the first strand cDNA (17, 18). In addition to these technical difficulties associated with RACE approaches, in many cases it is often necessary to repeat this process numerous times to obtain a full protein coding sequence (13).

25 To overcome the problem of fragmented cloning and inefficient second strand cDNA priming, procedures have been described that employ PCR-selection of fully transcribed cDNA by ligation of defined oligo ribonucleotides to the de-

5

10

25

5

10

15

Example 1: The Complete Open Reading Frame (C-ORF)
Technology: A Simple and Efficient Approach for Obtaining
the Entire Protein Coding Region of Genes

Cell cultures: HO-1 human melanoma cells (32) were grown in DMEM supplemented with 10% fetal bovine serum, 100 U/ml penicillin G and 100 μ g/ml streptomycin in a 5% CO₂ 95% air humidified incubator at 37°C. Du-145, a human prostate carcinoma cell line, was cultured in RPMI supplemented with the same reagents. Cells were maintained in the logarithmic phase of growth by subculturing 1:10 approximately two times per week.

C-ORF method for 5' cDNA end cloning: C-ORF consists of three reaction steps, reverse transcription (RT), second

strand synthesis and PCR. RNA samples (2 μ g) are reverse transcribed by SuperScript RT II (RNase H minus MMLV RT, BRL) with minor modifications from the manufacturer's protocol. The modifications include the use of 5 mM DTT instead of 10 mM DTT, 2 pmole gene specific RT primer for oligo dT and inclusion of 5 U RNaseIn. The RT reaction temperature was 45°C unless specified. First strand cDNA was purified with GlassMax(BRL) after RNase H (2.2 U) plus RNase A (0.5 μ g) digestion for 30 min at 37°C (50 μ l final).

10 A degenerate stem-and-loop annealing primer (D-SLAP or D-CLAP) was annealed in a 20 μ l mixture of 10 to 16 μ l cleaned first strand cDNA, 2 pmole annealing primer and 2 μ l 10X KlenTaq™ buffer (0.4 M Tricine-KOH, pH 9.2, 0.15 M KOAc, 35 mM Mg (OAc)₂ and 37.5 μ g/ml BSA). The annealing mixture was

15 incubated at 95°C for 1 min and was gradually cooled at 5°C/min to annealing temperature (42°C unless specified). During the 5 min of incubation at the annealing temperature, the annealing mixture was supplemented with 5 μ l of a polymerase mixture consisting of 0.25 μ l Advantage cDNA

20 polymerase mix™, 0.5 μ l 10 mM dNTPs and 0.5 μ l 10X KlenTaq™ buffer and incubated for 30 min at 68°C. Temperature was controlled using an MJ Minicycler™. Primary PCR was performed in a 25 μ l reaction consisting of 5 μ l of a second strand synthesis reaction mixture, 2.0 μ l 10X KlenTaq™

25 buffer, 200 μ M dNTPs, 5.0 pmole 3' gene specific primer (GSP), 10 pmole anchor primer and 0.25 μ l Advantage cDNA polymerase mix™. Basic PCR parameter, which varied depending on target size, were as follows: 95°C for 1 min, 27 cycles of amplification at 95°C for 30 sec, 58°C for 1

min and 68°C for a specified period (1 to 1.5 min per 1 kb target size), and 5 or 10 min further incubation at 68°C. For nested PCR, 0.5 μ l of the primary PCR was used employing essentially the same PCR parameters. Nested PCR reactions
5 (50 μ l) contain 0.5 μ l of primary PCR reaction mixture, 5 μ l 10X KlenTaq™ buffer, 0.2 mM dNTPs, 10 pmole nested GSP, 10 pmole anchor primer and 0.5 μ l Advantage cDNA polymerase mix™. A single primer reaction with GSP only or the anchor primer only is also performed with primary PCR reactions to
10 distinguish C-ORF artifacts. PCR reactions are resolved in 1% agarose gels and bands are purified with a gel purification kit (Qiagen). Purified bands are directly sequenced with anchor primer and GSP.

15 ***C-ORF method for 3' cDNA end cloning:*** Reverse transcription is performed and the product is purified as described in the C-ORF method for 5' end cloning in the previous section except that 10 pmole of the T-SLAP reagent and 5 μ g total RNA are used in place of the GSP. The primary PCR reaction
20 mixture (50 μ l) contains 2 μ l of purified first strand cDNA, 5 μ l 10X KlenTaq™ buffer, 0.2 mM dNTPs, 10 pmole GSP, 10 pmole anchor primer and 0.5 μ l Advantage cDNA polymerase mix™. Nested PCR mixtures consist of the same components except 1 μ l of a 10-fold diluted primary PCR of the first
25 strand cDNA synthesis and substitution of nested GSP for GSP. PCR parameters including temperature and duration are the same as described in the previous section.

Northern and Southern blot hybridizations: RNA samples

separated in 2%-formaldehyde agarose gels were transferred to Nylon membranes. Northern blots were hybridized with ³²P-labeled mda-5 3' EST (0.4 kb) or a gel purified C-ORF product (1.8 kb) as previously described (2). A Southern blot was prepared by transferring the PCR samples resolved in a 1% agarose gel to the Nylon membrane after depurination and denaturation. The blot was hybridized with ³²P-labeled nested GSP (M5R4) (Table 1) in 50% formamide hybridization buffer at room temperature.

RESULTS AND DISCUSSION

The C-ORF method was designed to obviate the complicated and inefficient steps associated with the conventional RACE procedure, which includes TdT-mediated tailing or single strand oligonucleotide ligation by T4 RNA ligase (16-18, 36). Figure 1A provides a schematic representation of the C-ORF strategy in which a universal primer site is generated by annealing the D-SLAP reagent to reverse transcribed cDNA during second strand cDNA synthesis. The hairpin (D-SLAP) or clover-leaf (D-CLAP) structure (Figure 1B) which form bulky loop structures are hypothesized to prevent the degenerate sequences from annealing and extending in the middle of reverse transcribed cDNA because of steric hindrance. Several restriction sites (Spe I, Xho I, Hinc II) for cloning PCR products into vectors are included in this construct. To ensure formation of the D-SLAP structure prior to association with target cDNA, the length of the stem is made longer than the degenerate sequences (18 vs. 12

nts) and the temperature from denaturation to annealing is gradually decreased (5°C/min) during second strand cDNA synthesis. In order to reduce PCR-related mutation and to enhance long range PCR amplification and specificity, the advantage cDNA polymeraseTM mixture (ClonTech, mixture of KlenTaq-1 DNA polymerase, proofreading Deep Vent_RTM and TaqStartTM antibody) is used during second strand cDNA synthesis and subsequent PCR amplification (37, 38).

10 In an attempt to obtain a complete open reading frame for the novel gene *mda-5* both library screening, conventional RACE and cap-switching RACE approaches were tried. Even after repeated attempts, these approaches resulted in the cloning of only a 2-kb cDNA library product of *mda-5* lacking
15 the complete open reading frame of this gene. In contrast, when the C-ORF cloning approach was used with 2 µg of total RNA from HO-1 cells a complete open reading frame for *mda-5* (~3.6 kb) was obtained. Employing an RT temperature of 48°C, an extended *mda-5* fragment of 1.8 kb was produced by
20 the C-ORF scheme with a single round using the D-SLAP reagent (Figure 2A).

The effect of concentration of the D-SLAP reagent during second strand synthesis and the anchor primer in primary PCR
25 reaction on *mda-5* gene amplification using the C-ORF approach were evaluated (Figure 2B). The specificity of the PCR products produced using the C-ORF approach with the D-SLAP reagent was determined by Southern blot hybridization (Figure 2C). Specific products of the anticipated size were

To confirm the utility of the C-ORF approach with the D-SLAP reagent for cloning the entire open reading frames from ESTs, studies were performed using three previously cloned genes (ISG-56, 1.5 kb (27); *mda-9*, 2 kb (28)) and *mda-5* (3.6 kb). For comparison, standard RT-PCR was performed with each message with a 5' primer designed from a distal 5' sequence and a 3' primer used in nested PCR reactions with C-ORF. As shown in Figure 3A, each C-ORF reaction produced

a fragment of the predicted size. Although complex banding patterns were seen in the C-ORF reactions versus RT-PCR, distinguishing PCR artifacts was readily accomplished by simultaneously running an anchor-only PCR reaction without the requirement for Southern blot hybridization verification (15). C-ORF-derived PCR products of ISG-56 (1.5 kb) (27) and *mda-9* (2 kb) (28) contained the full ORF for these genes, but they lacked 34 nts (ISG-56) and 27 nts (*mda-9*), respectively, from the complete sequence reported in GenBank. The C-ORF approach with the D-SLAP reagent using *mda-5* sequence information generated 2 fragments, a 3 kb product that was identical to that obtained previously and a 1.2 kb fragment representing a prematurely terminated gene product (31). For practical purposes, to obtain the most sequence information and the complete ORF for a given EST, the larger sized fragment would be the optimal fragment to isolate and sequence when performing the C-ORF approach. The appropriate size of the cDNA can also be obtained by Northern blotting analysis using the EST as a probe.

The C-ORF approach with the D-SLAP reagent has been used successfully for obtaining the complete ORF for a number of additional genes. Applications of the C-ORF approach using the D-SLAP reagent with the prostate carcinoma tumor antigen-1 (PCTA-1) gene (29) has resulted in the identification of variants of this gene with sizes of 3.5, 3.2, and 3.0 kb. The C-ORF approach permitted an extension of the 5' UTR sequence reported for PCTA-1 (29) by 400 bp. Additionally, the C-ORF approach with the D-SLAP reagent was

used to determine the sequence of a cDNA associated with cellular senescence and terminal cell differentiation, OLD-35 (34). A number of novel ESTs have been identified using an approach called reciprocal subtraction differential RNA display (RSDD) (33). These include novel ESTs that display elevated expression as a function of cancer progression, called progression elevated genes (PEGen), or reduced expression during cancer progression, called progression suppressed genes (PSGen) (33). C-ORF with the D-SLAP reagent has been used to generate a complete open reading frame for PEGen 28 (1.6 kb), PEGen 42 (1.8 kb) and PSGen 12 (1.2 kb).

Although the reaction temperature of RT profoundly affects the efficiency of C-ORF for genes like *mda-5*, a high RT temperature is not mandatory for other C-ORF applications (RT can be performed at 42 to 45°C). RT temperature-dependence may be a consequence of extensive secondary structure at the 5' UTR of specific cDNAs (6). An initial application of the C-ORF approach for cloning OLD-35 produced a shorter than anticipated product where a strong hairpin structure is predicted (34). However, an additional round of C-ORF produced a full ORF of OLD-35. Numerous attempts to clone OLD-35 by conventional RACE only yielded a few hundred-bp PCR product and further extension using different primer sets was not possible. In general, increasing RT temperature should prove helpful in assuring first strand cDNA synthesis to the end of the transcript by overcoming secondary structure of the RNA (21). By using

RNase H-minus MMLV-RT (Superscript II), reverse transcription can be efficiently performed up to 48°C and even higher temperatures (55 to 65°C) can be achieved using commercially available thermostable RT (14,21).

5

Second strand cDNA synthesis appears to depend on the D-SLAP concentration with lower concentrations of the D-SLAP reagent being less effective in promoting second strand cDNA synthesis (Figure 2B lane 2 and 3). Additional experiments suggest that the concentration of the D-SLAP reagent should be higher than 20 nM. Combinations of different anchor-gene specific primer (GSP) concentration ratios in the primary PCR were also tested for ISG-56. Combinations of 0.2 or 0.4 μ M anchor plus 0.2 μ M GSP produced an appropriate sized PCR product. However, the yield of PCR product was generally higher when using concentrations of 0.4 μ M anchor and 0.2 μ M GSP. Since 0.4 pmole of the D-SLAP reagent (1/5 of the second strand cDNA synthesis reaction) is carried over to the primary PCR reaction and it contains anchor primer binding sites, it may be necessary to use additional anchor primers to obtain appropriate PCR amplification. Thus, it is possible that the inefficient second strand cDNA synthesis at low D-SLAP and high anchor primer concentrations can result in a high yield of non-specific PCR products in *mda-5* (Figure 1B, lane 3).

10
15
20
25

Parameters for D-SLAP reagent annealing during second strand cDNA synthesis, including annealing temperature, temperature ramping and single tube second strand cDNA synthesis

003217 8204260

followed by PCR, were further investigated with *mda-9* (28) and ISG-56 (27). It was found that annealing temperatures up to 50°C were as effective as 42°C, and in certain cases superior to the lower temperature. Moreover, as observed
5 with ISG-56 (27), an annealing temperature higher than 46°C yielded a less complex pattern of C-ORF products. The annealing temperature used for C-ORF is considerably higher than the calculated T_m (44.5°C) for the D-SLAP priming site sequence of ISG-56 (27). It appears that raising the
10 annealing temperature prevents priming of D-SLAP reagents to relatively weak internal sites. A similar temperature dependence was also observed in single tube second strand cDNA synthesis and PCR, but the overall production of the appropriate PCR product in the single tube reaction was
15 significantly lower than that of the standard C-ORF protocol. This probably results because the D-SLAP reagent containing the anchor primer site may interfere with subsequent PCR processivity. Rapid ramping to 85°C followed by a gradual decrease to the annealing temperature (42°C,
20 5°C/min) or the addition of pre-annealed D-SLAP reagent did not significantly enhance PCR yield as compared with the standard C-ORF protocol.

The results described above clearly demonstrate the
25 efficiency of C-ORF with the D-SLAP reagent in cloning the open reading frame of a cDNA using EST information. However, although C-ORF effectively extends cDNAs to include the complete ORF, it did not extend the product to the end of the transcript. Provided that the stem and loop

structure of the D-SLAP reagent prevents degenerate sequences from binding to internal sites as predicted, it is possible that the RT did not extend to the 5' end of the transcript. Alternatively, the D-SLAP reagent may bind to internal sites during second strand cDNA synthesis resulting in the generation of a shorter product. The 5' end of sequences of ten C-ORF products where the degenerate region of D-SLAP reagent primed are summarized in Table 2. Considering that the G/C content of the 12 bp 5' sequences are 71% on average or 100% for *mda-5* (31) and *mda-9* (28), respectively, it appears that the annealing of the D-SLAP reagent is not completely sequence-independent but rather prefers G/C rich regions. In fact, C-ORF PCR yields are higher for *mda-9* (28) than for ISG-56 (27), although the RT-PCR yields of the two products are similar. Furthermore, while ISG-56 (27) C-ORF largely depends on which second strand cDNA synthesis procedure is used, *mda-9* (28) is easily amplified by most of the protocols tested.

Current data suggests that the target sequence used in C-ORF is important, but it is not the most critical determinant for annealing of the D-SLAP reagent to target sequences. Sequence analysis of 12 bp stretches in *mda-9* reveals one G/C site of 12/12, one G/C site of 11/12, one G/C site of 10/12 and three G/C sites of 9/12 (28). The C-ORF product of *mda-9* begins at the highest G/C content site that is located 27 bp downstream of the reported cDNA end of this cDNA. Although the D-SLAP reagent annealed to the highest G/C rich site in *mda-9*, based on the sequence analysis of

annealing sites (Table 2) this primer could also have annealed to other sites with > 75% G/C content in *mda-9*. In contrast, the D-SLAP reagent annealed to ISG-56 at the 58.3% G/C content site, whereas ISG-56 has two 12 bp stretches of 10/12 G/C and 6 sites of 9/12 G/C, respectively (27). If the G/C sequence content were the primary determinant of D-SLAP annealing and assuming a preference bias of PCR for shorter fragments, C-ORF applied to ISG-56 would be predicted to generate a mixture of products including the annealing site containing the 58.3% G/C content. The D-SLAP reagent could potentially anneal to all of the G/C rich sites, but the C-ORF product of ISG-56 that was amplified by PCR derived from a site of lower G/C content (58.3%) than the average G/C content of D-SLAP annealing sites (71%).

The propensity of C-ORF to generate near-end products may result either from preferential second strand cDNA synthesis when the D-SLAP reagent is annealed at the 3' end of the first strand cDNA or from an unanticipated PCR bias for the annealed D-SLAP reagent. In either case, the stem-and-loop structure of D-SLAP certainly plays a significant role in the preferential production of near-end fragments when using the C-ORF method. This bias may result by structurally preventing efficient second strand cDNA synthesis from deep internal sites within the cDNA. By providing a rigid stem-and-loop structure at the annealing temperature, it is possible that the longer stretch of DNA in the displaced strand that results from D-SLAP annealing has a greater chance to interfere with second strand cDNA synthesis. This

5 It is assumed that the stem-and-loop structure of D-SLAP facilitates second strand cDNA synthesis from the end or near the end of the cDNA. In this context, degenerate clover-leaf annealing primers (D-CLAP), containing three stem-and-loop structures, might even be more effective than the D-SLAP reagent in inhibiting second strand cDNA synthesis from internal priming sites, thereby alleviating the G/C sequence-dependence of D-SLAP. It is also established that RT has TdT-like activity and can add multiple Cs at the 3' end of the first strand cDNA from capped RNA (10,11). Based on these considerations, we designed two D-CLAP reagent, D-CLAP1 and D-CLAP2. While the D-CLAP1 reagent contains an annealing site with the sequence GGGN₁₀, the D-CLAP2 reagent contains 13 random oligonucleotides. C-ORF with the D-CLAP1 and D-CLAP2 reagents was performed with ISG-56 (27), *mda-9* (28), *mda-5*, PCTA-1 from either poly A site (3.5 and 5 kb, respectively (29, 39) and fibronectin (8 kb) (30). A single round of the C-ORF approach performed with the D-CLAP1 reagent generated bands of the expected size, not only for shorter transcripts (ISG-56 and *mda-9*) but also for larger transcripts (PCTA-1/pA and fibronectin) (Figure 4A), although this approach did increase band complexity. The annealing site sequence of ISG-56 is further upstream of the C-ORF product performed using the D-SLAP reagent, but it is still shorter than the

directly sequence the D-CLAP1 reagent C-ORF product with the anchor primer were unsuccessful, probably because of the hairpin structure in the anchor primer and the heterogeneous annealing of the GGG stretch, e.g., GGG can anneal in either direction to CCCC or CCCC. Provided that C-ORF with the D-CLAP2 reagent can yield the full protein coding region as exemplified in a number of applications, C-ORF using the D-CLAP2 reagent would be preferable to performing C-ORF with the D-CLAP1 reagent.

10

The utility of the D-SLAP reagent and C-ORF in cloning the 3' end of a cDNA was investigated to determine if this method could be used to obtain a complete 3' UTR of the PCTA-1 cDNA (29). An oligo dT stem and loop annealing primer (T-SLAP) was made by replacing oligo-dT for the random sequence of the D-SLAP reagent and was included in an RT reaction during the C-ORF procedure. PCR amplification (primary and nested) resulted in a specific 1.2-kb fragment, which was confirmed by genomic sequence and RT-PCR as PCTA-1 (Figure 5). The same protocol was also applied to 3' end cloning of OLD-35 (34), PEGen 28 and PEGen 50 (33) resulting in a 0.5 to 1 kb PCR amplification product corresponding to the 3' region of these cDNAs. These results indicate that primers based on a stem-and-loop structure can be efficiently used for both 5' and 3' end cloning of cDNAs.

25

Although the C-ORF method employs a common strategy involving a single side specific PCR reaction that is also used in the conventional RACE approach (14), C-ORF is unique

003317 0204260

in the generation of a universal anchor primer site and in the second strand cDNA synthesis step. Instead of homopolymer-tailing by TdT (15, 16) or single strand oligonucleotide ligation by T4 RNA ligase to the first strand cDNA (17,18), C-ORF involves the annealing of the D-SLAP or D-CLAP reagent to the first strand cDNA. By annealing degenerate primers, the C-ORF method eliminates an inherent problem of the conventional RACE approach caused by the inefficiency and sequence-dependence of the TdT and T4 RNA ligase reactions and the purification procedures required after the reaction (13, 23). Furthermore, since Taq polymerase is used during second strand cDNA synthesis and in the subsequent PCR reaction, no additional enzymes are required for the reaction. The complicated ramping to annealing temperature should not pose a problem, since most commercial thermocyclers have a program for gradual temperature declination. Compared with the conventional RACE approach, the C-ORF protocol represents a significant improvement resulting in less band complexity and a dramatic increase in the size of the cloned PCR generated product (13, 16-18, 20.) Although the anchor primer can on occasion generate PCR artifacts, it is easy to identify these spurious products by running anchor primer samples side by side with anchor primer plus GSP samples. This eliminates the need for Southern blot hybridization to confirm the gene specific authenticity of the PCR amplified product (15). A prominent feature of C-ORF that distinguishes it from the conventional and modified RACE approaches is that C-ORF yielded full protein coding regions in a single application

in 9 out of 10 test cases. Targets as small as 0.3 kb and as large as 8 kb could be amplified by the C-ORF protocol in a single reaction. These results confirm that the C-ORF approach will have wide applicability for identifying the complete protein coding regions of most ESTs and in many cases this will be achieved by a single application of the C-ORF methodology.

Applications of the C-ORF approach for cloning the complete open reading frame of diverse genes demonstrates that this approach is an efficient and simple way of cloning both 3' and 5' ends of cDNAs from EST sequences. With a small amount of starting material (1/5 to 1/3 of an RT reaction with 2 μ g of total RNA), C-ORF efficiently amplifies the complete protein coding region of diverse genes in most cases without the need for repetitive applications. Although in its present form C-ORF does not appear applicable for identifying the transcription start site, C-ORF is a cost-, time- and laborsaving method over currently used methods such as library screening and RACE for quickly obtaining biologically meaningful sequence data produced by C-ORF permits a determination of the uniqueness of the reported ESTs and provides sufficient information to procure corresponding clones from various providers to rapidly determine the full-length sequence of the gene. The quickness, ease of performance and efficiency of the C-ORF approach enables one to pursue multiple cDNA cloning projects simultaneously, which is necessary to expedite ongoing genome projects.

Table 1

Table 1. Sequence of gene specific primers used in the C-ORF cloning technology.

Primer	Gene	Sequence
AP-1	D-SLAP	5' TTCTGGTCGACTAGTGTTTAAACTCGAGAC 3'
AP-2	D-CLAP	5' CACGATCAGTCGACGAAAGTCGACACTCGAG 3'
M5R1	mda-5	5' TTTT TTTT TTTT CAGAGTAAACAATC 3'
M5R2	mda-5	5' TGTGCACCATCATTTGTTCCCAAGCC 3'
M5R3	mda-5	5' AATCTGACATTGGACTCATTTGAC 3'
M5R4	mda-5	5' GTTGAATCCTTGTTCATTATTCTAG 3'
M5S1	mda-5	5' GCCTGAGAGCCCTGTGGACAACCTCG 3'
56R1	ISG-56	5' GTGGCTGATATCTGGGTGCCCTAAGG 3'
56R2	ISG-56	5' CCTAAGGACCTTGTCTCACAGAGTTC 3'
56S1	ISG-56	5' CCAGATCTCAGAGGAGCCTGGCTAAGC 3'
M9R1	mda-9	5' AATCAGGATAAAGTGTCAACTATC 3'
M9R2	mda-9	5' ATCCCAAAGTAGCTAGGTTACATAATC 3'
M9S1	mda-9	5' CCTCAGAAGTCCGTGCCAGTGACCCGG 3'
FnR1	fibronectin	5' TTTT TTTT TTTT TGTGGAATGTAAATC 3'
FnR2	fibronectin	5' AGATGGATCTTGGCAGAGAGACATGC 3'
PCTAR1	PCTA-1	5' GAAGAAGTAGAACATCAGTGCC 3'
PCTAR2	PCTA-1	5' TCTTCTGTACAGCAGTATCTTACAT 3'
PCTAR3	PCTA-1	5' TTTT TTTT TTTT TTTTGTTCATGCGG 3'
PCTAR4	PCTA-1	5' TTACAAACAGCTCCCAAAATGTTGAAACT 3'

Table 2

Table 2. Sequence of the 5' end of the C-ORF products.

Name	5' sequence	G/C score	% G/C
<i>mda-5</i>	GCGCGCCGGC CT	11/12	91.7
ISG-56	TGCAGAACGG CT	7/12	58.3
<i>mda-9</i>	GGCGGCGGCG GC	12/12	100.0
PCTA-1A	TGGAGGCCTG GA	8/12	66.7
PCTA-1B	GCCAGTGCCT CA	8/12	66.7
PCTA-1C	CGATGTGGCC TT	7/12	58.3
OLD-35	CGGAGGACCA AT	7/12	58.3
PSGen 12	GCGGTGGTGA CG	9/12	75.0
PEGen 28	GTGTGGTGTG TC	7/12	58.3
PEGen 42	GGCGTTGCGA CG	9/12	75.0
G/C score	8994867986 74	85/120	70.8

09440941200

- 1999-2014, 1988.
24. Carninci, P., Kvam, C., Kitamura, A., Ohsumi, T.,
Okazaki, Y., Itoh, M., Kamiya, M., Shibata, K., Sasaki,
N., Izawa, M., Muramatsu, M., Hayashizaki, Y., and
5 Schneider, C. High-efficiency full-length cDNA cloning
by biotinylated CAP trapper, *Genomics*. 37: 327-336,
1996.
25. Kato, S., Sekine, S., Oh, S.W., Kim, N.S., Umezawa, Y.,
Abe, N., Yokoyama-Kobayashi, M., and Aoki, T.
10 Construction of a human full-length cDNA bank, *Gene*.
150: 243-250, 1994.
26. Hirzmann, J., Luo, D., Hahnen, J., and Hobom, G.
Determination of messenger RNA 5'-ends by reverse
transcription of the cap structure, *Nucleic Acids Res.*
15 21: 3597-3598, 1993.
27. Bahring, S., Sandig, V., Lieber, A., and Strauss, M.
Mapping of transcriptional start and capping points by
a modified 5' RACE technique, *Biotechniques*. 16: 807-
808, 1994.
- 20 28. CapFinder PCR cDNA Synthesis Kit, CLONTECHniques. 11:
2-4, 1996.
29. Hooft van Huijsduijnen, R. PCR-assisted cDNA cloning:
a guided tour of the minefield, *Biotechniques*. 24: 390-
392, 1998.
- 25 30. Schaefer, B.C. Revolutions in rapid amplification of
cDNA ends: new strategies for polymerase chain reaction
cloning of full-length cDNA ends, *Anal Biochem*. 227:
255-273, 1995.
31. Frohman, M.A., Dush, M.K., and Martin, G.R. Rapid

- 5 32. Loh, E.Y., Elliot, J.F., Cwirla, S., Lanier, L.L., and
Davis, M.M. Polymerase chain reaction with single-sided
specificity: analysis of T cell receptor delta chain,
Science. 243: 217-220, 1989.
- 10 33. Troutt, A.B., McHeyzer-Williams, M.G., Pulendran, B.,
and Nossal, G.J. Ligation-anchored PCR: a simple
amplification technique with single-sided specificity,
Proc Natl Acad Sci U S A. 89: 9823-9825, 1992.
- 15 34. Edwards, J.B., Delort, J., and Mallet, J.
Oligodeoxyribonucleotide ligation to single-standed
cDNAs: a new tool for cloning 5' ends of mRNA and for
constructing cDNA libraries by in vitro amplification,
Nucleic Acids Res. 19: 5227-5232, 1991.
- 20 35. Liu, X. And Gorovsky, M.A. Mapping the 5' and 3' ends
of Tetrahymena thermophila mRNAs using RNA ligase
mediated amplification of cDNA ends (RLM-RACE), Nucleic
Acids Res. 21: 4954-4960, 1993.
- 25 36. Frohman, M. RACE: Rapid Amplification of cDNA Ends. In:
M. Innis, D. Gelfand, J. Sninsky, and T. White (eds.),
PCR Protocols, pp. 28-38. San Diego, CA: Academic
Press, 1990.
37. Carninci, P., Nishiyama, Y., Westover, A., Itoh, M.,
Nagaoka, S., Sasaki, N., Okazaki, Y., Muramatsu, M.,
and Hayashizaki, Y. Thermostabilization and
thermoactivation of thermolabile enzymes by trehalose

and its application for the synthesis of full length cDNA, Proc Natl Acad Sci U S A. 95: 520-524, 1998.

38. Rychlik, W. In: B. White (ed.) Methods in Molecular Biology, Vol. 15, pp. 31-40. Totowa, NJ: Humana Press, 1993.
39. Chenchik, A., Diachenko, L., Mogadam, F., Tarabykin, V., Lukyanov, S., and Siebert, P.D. Full-length cDNA cloning and determination of mRNA 5' and 3' ends by amplification of adaptor-ligated cDNA, Biotechniques. 21: 526-534, 1996.
40. Maunders, M. DNA and RNA ligases. In: M. Burrell (ed.) Methods of Molecular Biology. Enzymes of Molecular Biology, Vol. 16, pp. 213-230. Totowa, NJ: Humana Press, 1993.
41. Maruyama, K. And Sugano, S. Oligo-capping: a simple method to replace the cap structure of eukaryotic mRNAs with oligoribonucleotides, Gene. 138: 171-174, 1994.
42. Fromont-Racine, m., Bertrand, E., Pictet, R., and Grange, T. A highly sensitive method for mapping the 5' termini of mRNAs, Nucleic Acids Res. 21: 1683-1684, 1993.
43. Wathelet, M., Moutschen, S., Defilippi, P., Cravador, A., Collet, M., Huez, G., and Content, J. Molecular cloning, full-length sequence and preliminary characterization of a 56-kDa protein induced by human interferons, Eur J Biochem. 155: 11-17, 1986.
44. Lin, J.J., Jiang, H., and Fisher, P.B. Melanoma differentiation associated gene-9, mda-9, is a human gamma interferon responsive gene, Gene. 207: 105-110,

5

- 10

15

What is claimed is:

1. A method for isolating a double-stranded cDNA having a nucleotide sequence of a complete open reading frame
5 which comprises:

A) admixing

- 10 (i) an isolated single-stranded cDNA,
- (ii) a first primer capable of forming a stem-loop structure, comprising
- 15 (a) at the 3' end of the primer, a first random sequence, linked to
- (b) a second sequence, linked to
- (c) a third sequence which forms a loop structure, linked to
- 20 (d) a fourth sequence, at the 5' end of the first primer, which is complementary to the second sequence,
- 25 under hybridization conditions sufficient for annealing the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and

(iii) a polymerase;

B) incubating the mixture from step (A) under suitable conditions for DNA synthesis; and

5

C) performing a polymerase chain reaction by admixing

(i) an aliquot of the mixture from (B),

10 (ii) a second primer which specifically binds to the single-stranded cDNA,

(iii) a third primer which comprises

15 (a) a fifth sequence identical to the third sequence of the first primer, linked to

(b) a sixth sequence identical to a portion of the second sequence of the first primer, and

20

(iv) a polymerase

25 under conditions suitable for a polymerase chain reaction so as to produce a double-stranded cDNA reaction product, thereby isolating the cDNA having the sequence of the complete open reading frame.

2. The method of claim 1, wherein the single-stranded DNA

09764029-11200

3. The method of claim 1, wherein the first primer has
5 the sequence
3'-
NNNNNNNNNNNNNCAGAGCTCAAATTTGTGATCAGCTGGTCTTTCACAAATTTGAGCTC
TG-5' (D-SLAP).

10 4. The method of claim 1, wherein the first primer has
the sequence
3'NNNNNNNNNNNGGGAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGTAGAAT
ACATCTTGAGCTAT-5' (D-CLAP1).

15 5. The method of claim 1, wherein the first primer has
the sequence
3'NNNNNNNNNNNNNNNAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGT
AGAATACATCTTGAGCTAT (D-CLAP2).

20 6. The method of claim 1, wherein the first primer
comprises an inosine nucleotide.

7. The method of claim 1, wherein the loop structure is a
simple loop structure, or a cloverleaf loop structure.

25 8. A method for generating a cDNA library which comprises:
A) admixing

(i) a population of single-stranded cDNA molecules which were reverse transcribed with an oligo-dT sequence linked to a defined nucleotide sequence,

5 (ii) a first primer capable of forming a stem-loop structure, comprising

(a) at the 3' end of the primer, a first random sequence linked to

10

(b) a second sequence, linked to

(c) a third sequence which forms a loop structure, linked to

15

(d) a fourth sequence, at the 5' end of the first primer, which is complementary to the second sequence,

20 under hybridization conditions sufficient for annealing the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and

25 (iii) a polymerase;

B) incubating the mixture from step (A) under suitable conditions for DNA synthesis by the polymerase; and

C) performing a polymerase chain reaction by admixing

(i) an aliquot of the mixture from (B),

5 (ii) a second primer which has the identical sequence
as the defined nucleotide sequence of the primer
in (A)(i),

(iii) a third primer which comprises

10

(a) a fifth sequence identical to the third
sequence of the first primer, linked to

15

(b) a sixth sequence identical to a portion
of the second sequence of the first
primer, and

(iv) a polymerase

20 under conditions suitable for a polymerase chain reaction so
as to produce double-stranded cDNA reaction products thereby
generating a cDNA library.

25 9. The method of claim 8, wherein the single-stranded DNA
is a cDNA reverse transcribed from an mRNA.

10. The method of claim 8, wherein the first primer has
the sequence 3'-
NNNNNNNNNNNNCAGAGCTCAAATTTGTGATCAGCTGGTCTTTCACAAATTTG

AGCTCTG-5' (D-SLAP) .

11. The method of claim 8, wherein the first primer has the sequence

5 3' NNNNNNNNNNGGAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGTAGAAT
ACATCTTGAGCTAT-5' (D-CLAP1).

12. The method of claim 8, wherein the first primer has the sequence

10 3' NNNNNNNNNNNNNNAGAGCTCACAGCTGAAGCAGCTGACTAGCACCTAGTGT
AGAATACATCTTGAGCTAT (D-CLAP2) .

13. The method of claim 8, wherein the first primer comprises an inosine nucleotide.

14. The method of claim 8, wherein the loop structure is a simple loop structure, or a cloverleaf loop structure.

15. A kit for the generation of a complete open reading
20 frame double-stranded cDNA of interest which comprises:

- (i) a first primer capable of forming a stem-loop structure, comprising

- 25 (a) at the 3' end of the primer, a first
random sequence linked to

- (b) a second sequence, linked to

- (c) a third sequence which forms a loop

structure, linked to

- 5 (d) a fourth sequence, at the 5' end of the first primer, which is complementary to the second sequence, and

- (ii) a second primer which comprises
- 10 (a) a fifth sequence identical to the third sequence of the first primer, linked to
- (b) a sixth sequence identical to a portion of the second sequence of the first primer.

16. A method for isolating a double-stranded cDNA having a nucleotide sequence of a complete open reading frame which comprises:
- 15

- (a) admixing
- 20 (i) a biological sample containing mRNA,
- (ii) a primer which forms a stem-loop structure, comprising:
- (a) a poly-T sequence at the 3' end of the primer linked to
- 25
- (b) a first random sequence linked to
- (c) a second sequence which forms a loop

structure linked to

5 (d) a third sequence at the 5' end of the primer
which is complementary to the first
sequence, and

(iii) a reverse transcriptase,

10 under hybridization conditions sufficient for annealing
the primer to the mRNA poly-A sequence;

(b) incubating the mixture from step (a) under suitable
conditions for reverse transcription;

15 (c) performing a polymerase chain reaction with an aliquot
of the mixture from step (b) using one gene-specific
primer which is pre-defined and one primer which has a
sequence identical to at least a portion of the primer
20 sequence of element (ii), thereby isolating the cDNA
having the sequence of the complete open reading frame.


17. The method of claim 16, wherein the primer has the
s e q u e n c e 3 ' -
TTTTTTTTTTTCAGAGCTCAAATTTGTGATCAGCTGGTCTTTCACAAATTTG
25 AGCTCTG-5' (T-SLAP).

Method for Full-Length cDNA Cloning Using Degenerate Stem-
Loop Annealing Primers

Abstract of the Disclosure

5 The present invention provides a method for isolating a double-stranded cDNA having a nucleotide sequence of a complete open reading frame which comprises: (A) admixing (i) an isolated single-stranded cDNA, (ii) a first primer capable of forming a stem-loop structure, comprising (a) at
10 the 3' end of the primer, a first random sequence, linked to (b) a second sequence, linked to (c) a third sequence which forms a loop structure, linked to (d) a fourth sequence, at the 5' end of the first primer, which is complementary to the second sequence, under hybridization conditions
15 sufficient for annealing the first sequence of the first primer to the sequence at the 3' end of the single-stranded cDNA, and (iii) a polymerase; (B) incubating the mixture from step (A) under suitable conditions for DNA synthesis; and (C) performing a polymerase chain reaction by admixing
20 (i) an aliquot of the mixture from (B), (ii) a second primer which specifically binds to the single-stranded cDNA, (iii) a third primer which comprises (a) a fifth sequence identical to the third sequence of the first primer, linked to (b) a sixth sequence identical to a portion of the second sequence
25 of the first primer, and (iv) a polymerase, under conditions suitable for a polymerase chain reaction so as to produce a double-stranded cDNA reaction product, thereby isolating the cDNA having the sequence of the complete open reading frame.


Figure 1A




Figures 1B and 1C

B.


D-SLAP



D-CLAP1




D-CLAP2




C.

T-SLAP




Figures 2A, 2B, and 2C




Figures 2D and 2E

D.




Lane 1- ctl with EST
Lane 2- IFN- β with EST
Lane 3- ctl with C-ORF product
Lane 4- IFN- β with C-ORF product

E.




Lane 1- 1 kb ladder
Lane 2- RT-PCR R2/S8
Lane 3- RT-PCR R4/S8

Figure 3




Figures 4A and 4B

A.




Lane 1 & 6- 1 kb ladder
 Lane 2- mda-9
 Lane 3- fibronectin
 Lane 4- PCTA-1 (R1)
 Lane 5- PCTA-1 (pA)
 Lane 7- mda-5
 Lane 8- ISG-56

B.



Lane 1- 1 kb ladder
 Lane 2- fibronectin
 Lane 3- PCTA-1 (pA)
 Lane 4- mda-9
 Lane 5- ISG 56

Figure 5



Lane 1- 1 kb ladder

Lane 2- AP only

Lane 3- GSP/AP

Lane 4- GSP only

DECLARATION AND POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHOD FOR FULL-LENGTH cDNA CLONING USING DEGENERATE STEM-LOOP ANNEALING PRIMERS

the specification of which:
(check one)

☒ is attached hereto

☐ was filed on _____ as

Application Serial No. _____

and was amended _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119 (a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International Application which designated at least one country other than the United States, listed below. I have also identified below any foreign application for patent or inventor's certificate, or PCT International Application having a filing date before that of the earliest application from which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

<u>Number</u>	<u>Country</u>	<u>Filing Date</u>	<u>Yes</u>	<u>No</u>
N/A				

05724028-113600

Applicants: Paul B. Fisher and Dong-chul Kang

U.S. Serial No.: Not Yet Known

Filed: Herewith

Declaration and Power of Attorney

Page 2

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

<u>Provisional Application No.</u>	<u>Filing Date</u>	<u>Status</u>
N/A		

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States Application(s), or Section 365(c) of any PCT International Application(s) designating the United States listed below. Insofar as this application discloses and claims subject matter in addition to that disclosed in any such prior Application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56, which became available between the filing date(s) of such prior Application(s) and the national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
N/A		

And I hereby appoint

John P. White (Reg. No. 28,678); Christopher C. Dunham (Reg. No. 22,031); Norman H. Zivin (Reg. No. 25,385); Jay H. Maioli (Reg. No. 27,213); William E. Pelton (Reg. No. 25,702); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Wendy E. Miller (Reg. No. 35,615); Richard S. Milner (Reg. No. 33,970); Robert T. Maldonado (Reg. 38,232); Paul Teng (40,837); Richard F. Jaworski (Reg. No. 33,515); Elizabeth M. Wieckowski (Reg. No. 42,226); Pedro C. Fernandez (Reg. No. 41,741); Gary J. Gershik (Reg. No. 39,992); Jane M. Love (Reg. No. 42,812); Spencer H. Schneider (Reg. No. 45,923) and Raymond A. Diperna (Reg. No. 44,063).

and each of them, all c/o Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, New York 10036, my attorneys, each with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent, to transact all business in the Patent and Trademark Office connected therewith and to file any International Applications which are based thereon under the provisions of the Patent Cooperation Treaty.

